

ON NODAL LINES OF NEUMANN EIGENFUNCTIONS

Krzysztof Burdzy

Abstract. We present a new method for locating the nodal line of the second eigenfunction for the Neumann problem in a planar domain.

1. Introduction. This note is inspired by recent progress on the “hot spots” conjecture of J. Rauch originally proposed in 1974. The conjecture states that the maximum of the second Neumann eigenfunction in a Euclidean domain is attained on the boundary. This formulation is somewhat vague but the counterexamples given by Burdzy and Werner (1999) and Bass and Burdzy (2000) show that even the weakest version of the conjecture fails in some planar domains. The first positive results on the “hot spots” conjecture appeared in Kawohl (1985) and Bañuelos and Burdzy (1999). The last article contains a proof that the conjecture holds in all triangles with an obtuse angle. The theorem was generalized to “lip domains”, i.e., planar domains between two graphs of Lipschitz functions with the Lipschitz constant 1; this more general result was hinted at in Bañuelos and Burdzy (1999) and proved in Atar and Burdzy (2002). See Atar (2001), Jerison and Nadirashvili (2000) and Pascu (2002) for some other positive results.

There is currently a significant gap between the best positive results and the strongest counterexamples to the “hot spots” conjecture. It is widely believed that the conjecture holds in all convex domains in $\mathbf{R}^n$ for all $n \geq 2$. But the problem remains open even for triangles whose all angles are acute.

One of the key assumptions in some theorems of Bañuelos and Burdzy (1999), and all results of Jerison and Nadirashvili (2000) and Pascu (2002), is the symmetry of the domain with respect to a straight line. The assumption is only needed to show that the nodal line

Research partially supported by NSF grant DMS-0071486.

AMS mathematics subject classification (2000): 35P99, 60J65

Key words or phrases: nodal line, reflected Brownian motion, mirror coupling, eigenfunction, Neumann problem.

of the second Neumann eigenfunction lies on the line of symmetry. Then one can replace the original Neumann problem on the whole domain with the mixed Dirichlet-Neumann problem on a nodal subdomain. As it often happens in analysis, the Dirichlet problem (in this case, the mixed Dirichlet-Neumann problem) turns out to be considerably easier to deal with than the original Neumann problem. Hence, one would like to be able to find the location of the nodal line for the second Neumann eigenfunction in domains which are not necessarily symmetric. If this goal is achieved, some further progress on the “hot spots” conjecture may be expected.

Very few methods for finding the nodal line are known; the following short list is probably complete.

- (a) If a domain has a line of symmetry, the nodal line lies on that line, subject to some natural extra assumptions.
- (b) There are a handful of classes of domains, such as rectangles and ellipses, for which explicit formulae for eigenfunctions are known.
- (c) One can find an approximate location of the nodal line in domains which are “long and thin” in the asymptotic sense; see Jerison (2000).

The purpose of this note is to add the following item to this list.

- (d) In some domains, one can use the probabilistic method of “mirror couplings” to delineate a region which the nodal line must intersect.

It should be emphasized that one cannot expect to find the location of the nodal line of the second Neumann eigenfunction in the sense of an explicit formula, except in some trivial cases. What one can realistically hope to achieve is to obtain sufficiently accurate information about the nodal line so that this information can in turn be used to prove some other results of interest; the combination of Lemmas 2 and 3 of Burdzy and Werner (1999) is an example of such a result on nodal lines.

Finally, we would like to mention the results of Melas (1992) on the nodal lines in the Dirichlet case.

2. Mirror couplings and nodal lines. Suppose that D is a planar domain (open and connected set) with a piecewise smooth boundary. Informally speaking, a mirror coupling is a pair (X_t, Y_t) of reflecting Brownian motions in $\overline{D}$, such that the line of symmetry K_t between X_t and Y_t does not change on any interval (s, u) such that $X_t \notin \partial D$ and $Y_t \notin \partial D$ for all $t \in (s, u)$. More formally, (X_t, Y_t) satisfy the following system of stochastic

differential equations,

$$dX = dW + dL, \quad dY = dZ + dM, \quad dZ = dW - 2m \cdot m \cdot dW, \quad m = \frac{Y - X}{|Y - X|},$$

for times less than $\zeta = \inf\{s : X_s = Y_s\}$. We set $Y_t = X_t$ for $t \geq \zeta$. Here W is a planar Brownian motion, and Z is another Brownian motion for which the increments are mirror images of those of W , the mirror being the line with respect to which X and Y are symmetric. The processes X and Y are reflecting Brownian motions with the singular drifts L and M on the boundary, i.e., $\int_0^\infty \mathbf{1}_{\partial D}(X_s) |dL_s| = 0$, and similarly for M . A rigorous construction of the mirror coupling was given in Atar and Burdzy (2002) although mirror couplings had been informally used in the past; see Atar and Burdzy (2002) for a short review of the history of this concept.

Recall that the first eigenvalue for the Laplacian in D with Neumann boundary conditions is equal to 0 and the corresponding eigenfunction is constant. The second eigenvalue need not be simple but its multiplicity can be only 1 or 2 (Nadirashvili (1986, 1988)). The nodal set is the set of points in D where the second eigenfunction vanishes. If D is not simply connected, the nodal set need not be a connected curve. The nodal set of any second Neumann eigenfunction divides the domain D into exactly two nodal domains (connected and open sets).

Theorem 1. *Suppose that $A \subset \overline{D}$ is closed, $x \in D \setminus A$, and D_1 is the connected component of $D \setminus A$ which contains x . Assume that for some $y \in D \setminus D_1$, the mirror K_t for the coupling starting from $(X_0, Y_0) = (x, y)$ satisfies $K_t \cap D \subset A$ for all $t \in [0, \zeta]$ a.s. Then for any second Neumann eigenfunction in D , none of its nodal domains can have a closure which is a subset of $\overline{D} \setminus A$ containing x .*

Proof. Suppose that B is one of the nodal domains for a second eigenfunction and $x \in \overline{B} \subset \overline{D} \setminus A$. Let C_1 be a non-empty open disc centered at a boundary point of B , so small that $B_* = B \cup C_1$ satisfies $x \in \overline{B_*} \subset \overline{D} \setminus A$, and there exists a non-empty open disc $C_2 \subset D \setminus B_*$. Assume that for some $y \in D \setminus D_1$, if the mirror coupling starts from $(X_0, Y_0) = (x, y)$ then $K_t \cap D \subset A$ for all $t \in [0, \zeta]$ a.s. Since the stationary measure for X_t is the uniform distribution in D , X_t will hit C_2 with probability one, and so it will leave B_* with probability 1. Recall that there are exactly two nodal domains— B is one of them; let B_1 denote the other one. Let $\tau_X(B_*) = \inf\{t \geq 0 : X_t \notin B_*\}$ and $\tau_Y(B_1) = \inf\{t \geq 0 : Y_t \notin B_1\}$. We will argue that $\tau_X(B_*) < \tau_Y(B_1)$ a.s. Suppose otherwise. By the continuity of $t \rightarrow Y_t$, $Y_{\tau_Y(B_1)}$ belongs to $\overline{\partial B_1} \setminus \overline{\partial D}$, and so $Y_{\tau_Y(B_1)} \in \overline{B} \subset \overline{B_*}$. We have

assumed that $\tau_X(B_*) \geq \tau_Y(B_1)$, so $X_{\tau_Y(B_1)} \in \overline{B}_*$. Since the points $X_{\tau_Y(B_1)}$ and $Y_{\tau_Y(B_1)}$ belong to the connected set $\overline{B}_*$, their line of symmetry $K_{\tau_Y(B_1)}$ must also intersect $\overline{B}_*$. We have $K_t \cap D \subset A$ for all $t \in [0, \zeta]$ so $A \cap \overline{B}_* \neq \emptyset$, which is a contradiction. We conclude that $\tau_X(B_*) < \tau_Y(B_1)$ a.s.

Let $\mu_2 > 0$ denote the second Neumann eigenvalue. Then μ_2 is the first eigenvalue for the mixed problem in the nodal domain B , with the Neumann boundary conditions on ∂D and the Dirichlet boundary conditions on the nodal line. The fact that B_* is strictly larger than B easily implies that $\mu_2 > \mu_2^*$, where μ_2^* is the analogous mixed eigenvalue for B_* . Using the well known identification of Brownian motion density with the heat equation solution, we obtain from Proposition 2.1 of Bañuelos and Burdzy (1999) that

$$\lim_{t \rightarrow \infty} P(\tau_Y(B_1) > t) e^{\mu_2 t} \in (0, \infty).$$

Since $\tau_X(B_*) < \tau_Y(B_1)$ a.s.,

$$\lim_{t \rightarrow \infty} P(\tau_X(B_*) > t) e^{\mu_2 t} < \infty,$$

and so $\mu_2^* \geq \mu_2$, but this contradicts the fact that $\mu_2 > \mu_2^*$. Our initial assumption that $x \in \overline{B} \subset \overline{D} \setminus A$ for some nodal domain B must be false. $\square$

We will illustrate our main result with two examples. Before we do that, we recall a few crucial facts about mirror couplings from Burdzy and Kendall (2000) and Bañuelos and Burdzy (1999).

Suppose that D is a polygonal domain and I is a line segment contained in its boundary. Let J denote the straight line containing I and recall that K_t denotes the mirror line. Let H_t denote the “hinge,” i.e., the intersection of K_t and J . Note that H_t need not belong to ∂D . Suppose that for all t in $[t_1, t_2]$, the reflected Brownian motions X_t and Y_t do not reflect from any part of ∂D except I . Let α_t denote the smaller of the two angles formed by K_t and J . Then all possible movements of K_t have to satisfy the following condition.

- (M) The hinge does not move within the time interval $[t_1, t_2]$, i.e., $H_t = H_{t_1}$ for all $t \in [t_1, t_2]$. The angle α_t is a non-decreasing function of t on $[t_1, t_2]$.

A domain D with piecewise smooth boundary can be approximated by polynomial domains D_n . Mirror couplings in the approximating domains D_n converge weakly to a mirror coupling in D . One can deduce which motions of the mirror K_t in D are possible by analyzing all allowed movements of mirrors in D_n ’s and then passing to the limit. We

will not present the details of this limit theorem here. The readers who are concerned about full rigor should add an extra assumption to Example 2 that D is a polygonal domain.

Example 1. Suppose that D is an obtuse triangle with vertices C_1, C_2 and C_3 . Let C_3 be the vertex with an angle greater than $\pi/2$. Let $\overline{C_j C_k}$ denote the line segment with endpoints C_j and C_k , and let $d(C_j, C_k)$ denote the distance between these points. Let the points $C_4, C_{10} \in \overline{C_1 C_3}$, $C_6, C_{11} \in \overline{C_2 C_3}$, and $C_5, C_7, C_8, C_9 \in \overline{C_1 C_2}$ be such that $d(C_1, C_4) = d(C_4, C_3) = d(C_1, C_5)$, $d(C_2, C_6) = d(C_6, C_3) = d(C_2, C_7)$, and the following pairs of line segments are perpendicular: $\overline{C_4 C_8}$ and $\overline{C_2 C_3}$, $\overline{C_6 C_9}$ and $\overline{C_1 C_3}$, $\overline{C_5 C_{10}}$ and $\overline{C_1 C_3}$, $\overline{C_7 C_{11}}$ and $\overline{C_2 C_3}$. See Figure 1.




Figure 1.

Let A be the closed subset of $\overline{D}$ whose boundary consists of line segments $\overline{C_3 C_4}$, $\overline{C_3 C_6}$ and $\overline{C_5 C_7}$ and circular arcs $\widehat{C_4 C_5}$ and $\widehat{C_6 C_7}$. The arcs $\widehat{C_4 C_5}$ and $\widehat{C_6 C_7}$ are centered at C_1 and C_2 , respectively. Let A_1 be the closure of the union of pentagons with vertices $C_3 C_4 C_8 C_9 C_6$ and $C_3 C_{10} C_5 C_7 C_{11}$.

The second Neumann eigenvalue is simple in obtuse triangles by a theorem of Atar and Burdzy (2002).

We will show that

- (i) The nodal line for the second Neumann eigenfunction must intersect A .
- (ii) The nodal line lies within A_1 .

Let α_t denote the angle formed by the mirror K_t with $\overline{C_1 C_2}$. Let β_1 and β_2 denote the angles such that if $\alpha_t = \beta_1$ then K_t is perpendicular to $\overline{C_2 C_3}$ and when $\alpha_t = \beta_2$ then

K_t is perpendicular to $\overline{C_1C_3}$. An argument totally analogous to the proof of Theorem 3.1 of Bañuelos and Burdzy (1999) shows that if $\alpha_0 \in [\beta_1, \beta_2]$ then $\alpha_t \in [\beta_1, \beta_2]$ for all $t \geq 0$ a.s.

Consider any point $x \in D \setminus A$ to the left of the arc $\widehat{C_4C_5}$. Let $y \in \overline{D}$ be such that C_1, x and y lie on the same line and $d(C_1, y) = d(C_1, C_3)$. Consider a mirror coupling (X_t, Y_t) with $X_0 = x$ and $Y_0 = y$. We will show that $K_t \cap D \subset A$ for all $t \geq 0$.

We have $K_0 \cap D \subset A$, by the definition of y . Assume that it is not true that $K_t \cap D \subset A$ for all t . We will show that this assumption leads to a contradiction. It follows from this assumption that for some $\varepsilon > 0$ there exists t with $d(C_1, K_t) \leq d(C_1, C_3)/2 - \varepsilon$ or $d(C_2, K_t) \leq d(C_2, C_3)/2 - \varepsilon$. Let T_1 be the infimum of such t . Without loss of generality assume that $d(C_1, K_{T_1}) \leq d(C_1, C_3)/2 - \varepsilon$. Let T_2 be the last time t before T_1 when $d(C_1, K_t) \geq d(C_1, C_3)/2 - \varepsilon/2$. Note that $d(C_1, K_t) \leq d(C_1, C_3)/2 - \varepsilon/2$ for all $t \in [T_2, T_1]$. This implies that $d(C_1, X_t) \leq d(C_1, C_3)$ and $d(C_1, Y_t) \leq d(C_1, C_3)$ for all $t \in [T_2, T_1]$. Hence, the processes X_t and Y_t can reflect only on $\overline{C_1C_2}$ and $\overline{C_1C_3}$ on the time interval $[T_2, T_1]$. According to **(M)**, the distance from C_1 to K_t cannot decrease between times T_2 and T_1 , which contradicts the definition of these times.

Claim (i) now follows from Theorem 1. The second claim is a consequence of the first one and the bounds on the direction of the gradient of the second eigenfunction proved in Theorem 3.1 of Bañuelos and Burdzy (1999).

We will argue heuristically that it is impossible to obtain much sharper estimates for the nodal line location without imposing some extra assumptions on the obtuse triangle. To see this, consider an obtuse triangle with two angles very close to $\pi/2$ —one slightly less than the right angle and another one slightly larger than that (see Figure 2). Such a triangle has a shape very close to a thin circular sector. Let us assume that the diameter of the triangle is equal to 1. The nodal line in a circular sector is an arc with center at its vertex C . The nodal line distance from C is equal to $a_0/a_1 \approx 0.63$, where a_0 is the first positive zero of the Bessel function of order 0 and a_1 is the first positive zero of its derivative. This follows from known results on eigenfunctions in discs and estimates for Bessel function zeroes, see Bandle (1980), p. 92. Our methods yield 0.5 as the lower bound for the distance of the nodal line from C ; we see that this estimate cannot be improved beyond 0.63.




Figure 2.

We end this example with a conjecture which, if proved, would considerably strengthen claims (i) and (ii).

Conjecture 1. Suppose that D is a triangle with vertices C_1, C_2 and C_3 and that the angle at C_3 is greater than $\pi/2$. Let $C_{12} \in \overline{C_1 C_2}$ be the unique point such that $\overline{C_{12} C_3}$ is perpendicular to $\overline{C_1 C_2}$, and let D_1 be the larger of the two connected components of $D \setminus \overline{C_{12} C_3}$. Then the nodal line for the second Neumann eigenfunction is contained in D_1 .

Little is known about eigenfunctions in triangles different from equilateral; see Pinsky (1980, 1985) for that special case.

Example 2. Consider a convex domain D in the plane whose boundary contains line segments $\{(x_1, x_2) : -a \leq x_1 \leq a, x_2 = 0\}$ and $\{(x_1, x_2) : -a \leq x_1 \leq a, x_2 = 1\}$ for some $a > 0$. Suppose that the remaining parts of ∂D are contained in $\{(x_1, x_2) : -a - b \leq x_1 \leq -a, 0 \leq x_2 \leq 1\}$ and $\{(x_1, x_2) : a \leq x_1 \leq a + b, 0 \leq x_2 \leq 1\}$. We will assume that $b < a - 1/(4a)$. This means that we have to have $a > 1/2$. Let

$$A = \{(x_1, x_2) \in \overline{D} : -b - 1/(4a) \leq x_1 \leq b + 1/(4a)\},$$

$$A_1 = \{(x_1, x_2) \in \overline{D} : -b - 1/(4a) - 1 \leq x_1 \leq b + 1/(4a) + 1\}.$$

Figure 3 shows an example of D and the corresponding rectangle A .



Figure 3.

We will prove that

- (i) The nodal line for any second Neumann eigenfunction must intersect A .

We will say that D is a “lip domain” if its boundary consists of graphs of two Lipschitz functions with the Lipschitz constant 1. The domain depicted in Figure 3 is a lip domain. The second eigenvalue is simple in every lip domain, by a result of Atar and Burdzy (2002).

Our second claim is

- (ii) If D is a lip domain then the nodal line for the Neumann eigenfunction lies within A_1 .

Suppose that $x \in D \setminus A$ lies to the left of A . If $x = (x_1, x_2)$ for some $x_1 \geq -a$ then let $y = (-x_1, x_2)$. Otherwise let y be any point with the first coordinate greater than or equal to a . Let (X_t, Y_t) be a mirror coupling with $X_0 = x$ and $Y_0 = y$. Our goal is to show that $K_t \cap D \subset A$ for all $t \geq 0$.

It is elementary, and so it is left to the reader, to check that for any points $x = (x_1, x_2)$ and $y = (y_1, y_2)$ in D with $x_1 \leq -a$ and $y_1 \geq a$, the line of symmetry for x and y intersects D within A . Hence $K_0 \cap D \subset A$, by the choice of y . Next we follow the same line of thought as in Example 1. Suppose that the condition $K_t \cap D \subset A$ fails for some $t \geq 0$. We will show that this assumption leads to a contradiction. It follows from our assumption that for some $\varepsilon > 0$ there exists t and $(z_1, z_2) \in K_t \cap \overline{D}$ with $z_1 \leq -b - 1/(4a) - \varepsilon$ or $z_1 \geq b + 1/(4a) + \varepsilon$. We can assume without loss of generality that $z_1 \leq -b - 1/(4a) - \varepsilon$. Let T_1 be the infimum of t such that $K_t \cap \overline{D}$ intersects the line $\{(z_1, z_2) : z_1 = -b - 1/(4a) - \varepsilon\}$. Let T_2 be the supremum of $t \leq T_1$ such that $K_t \cap \overline{D}$ does not intersect $\{(z_1, z_2) : z_1 = -b - 1/(4a) - \varepsilon/2\}$. Write $X_t = (X_t^1, X_t^2)$ and $Y_t = (Y_t^1, Y_t^2)$. The following argument applies when $\varepsilon > 0$ is sufficiently small—obviously, we can make such an assumption on ε . Note that $X_t^1 \leq a$ and $Y_t^1 \leq a$ for $t \in [T_2, T_1]$. During this interval, Y_t can only reflect on the horizontal parts of the boundary of D , so according to **(M)**, this reflection cannot decrease $\inf\{z_1 : (z_1, z_2) \in K_t \cap \overline{D}\}$. The same holds true if X_t reflects on the horizontal parts of ∂D . Suppose that X_t reflects on the part of ∂D to the left of $\{(z_1, z_2) : z_1 = -a\}$ which is not horizontal. Then the hinge H_t lies outside $\overline{D}$ and it easily follows from **(M)** that both intersection points of K_t with ∂D move to the right. Hence, $\inf\{z_1 : (z_1, z_2) \in K_t \cap \overline{D}\}$ is a non-decreasing function of t for $t \in [T_2, T_1]$. This contradicts the definition of T_1 and T_2 .

The above argument and Theorem 1 yield claim (i). The second claim follows from the first one and the bounds on the direction of the gradient of the second eigenfunction given in Example 3.1 of Bañuelos and Burdzy (1999).

REFERENCES

- [1] R. Atar (2001). Invariant wedges for a two-point reflecting Brownian motion and the “hot spots” problem. *Elect. J. of Probab.* **6**, paper 18, 1–19.
- [2] R. Atar and K. Burdzy (2002), On Neumann eigenfunctions in lip domains (preprint).
- [3] C. Bandle *Isoperimetric Inequalities and Applications*. Monographs and Studies in Mathematics, 7. Pitman, Boston, Mass.-London, 1980.
- [4] R. Bañuelos and K. Burdzy (1999), On the “hot spots” conjecture of J. Rauch. *J. Func. Anal.* **164**, 1–33.
- [5] R. Bass and K. Burdzy (2000) Fiber Brownian motion and the ‘hot spots’ problem *Duke Math. J.* **105**, 25–58.
- [6] K. Burdzy and W. Kendall (2000) Efficient Markovian couplings: examples and counterexamples *Ann. Appl. Probab.* **10**, 362–409.
- [7] K. Burdzy and W. Werner (1999) A counterexample to the “hot spots” conjecture. *Ann. Math.* **149**, 309–317.
- [8] D. Jerison (2000) Locating the first nodal line in the Neumann problem. *Trans. Amer. Math. Soc.* **352**, 2301–2317.
- [9] D. Jerison and N. Nadirashvili (2000) The “hot spots” conjecture for domains with two axes of symmetry. *J. Amer. Math. Soc.* **13**, 741–772.
- [10] B. Kawohl, *Rearrangements and Convexity of Level Sets in PDE*, Lecture Notes in Mathematics 1150, Springer, Berlin, 1985.
- [11] A. Melas (1992) On the nodal line of the second eigenfunction of the Laplacian in R^2 . *J. Differential Geom.* **35**, 255–263.
- [12] N.S. Nadirashvili (1986) On the multiplicity of the eigenvalues of the Neumann problem, *Soviet Mathematics, Doklady*, **33**, 281–282.
- [13] N.S. Nadirashvili (1988) Multiple eigenvalues of the Laplace operator, *Mathematics of the USSR, Sbornik*, **133-134**, 225–238.
- [14] M. Pascu (2002) Scaling coupling of reflecting Brownian motions and the hot spots problem *Trans. Amer. Math. Soc.* (to appear)
- [15] M. Pinsky (1980) The eigenvalues of an equilateral triangle. *SIAM J. Math. Anal.* **11**, 819–827.
- [16] M. Pinsky (1985) Completeness of the eigenfunctions of the equilateral triangle. *SIAM J. Math. Anal.* **16**, 848–851.

Department of Mathematics
University of Washington
Box 354350
Seattle, WA 98195-4350, USA

Email: burdzy@math.washington.edu
<http://www.math.washington.edu/~burdzy/>